

COMMENTARY

10.1002/2017JA024679

Key Points:

- Different storms have different mixes of ring current strength, radiation-belt enhancement, plasma sheet temperature, auroral currents, etc
- The *Dst* index is a poor identifier of high-speed-stream-driven storms and of other types of geospace storms
- A call is made to the space physics community to refine the definitions of storms and to develop criteria to identify them

Correspondence to:

J. E. Borovsky,
jborovsky@space-science.org

Citation:

Borovsky, J. E., & Shprits, Y. Y. (2017). Is the *Dst* index sufficient to define all geospace storms? *Journal of Geophysical Research: Space Physics*, 122. <https://doi.org/10.1002/2017JA024679>

Received 14 AUG 2017

Accepted 7 OCT 2017

Accepted article online 16 OCT 2017

Is the *Dst* Index Sufficient to Define All Geospace Storms?

Joseph E. Borovsky¹ and Yuri Y. Shprits^{2,3}
¹Space Science Institute, Boulder, CO, USA, ²GFZ, German Research Center for Geosciences, Potsdam, Germany,

³Department of Earth and Space Sciences, UCLA, Los Angeles, CA, USA

Abstract The purpose of this commentary is (1) to raise awareness about some shortcomings of the use of the *Dst* index to identify storms, to gauge storm intensity, and to represent storm time space-weather phenomena and (2) to initiate discussions about different types of storms and about improved identifiers for different types of storms.

Dst, the *H*-component perturbation on equatorial magnetometers (Mayaud, 1980), is a historical definition of a magnetic storm that goes back to von Humboldt in the early 1800's (Malin & Barraclough, 1991). The creation of the *Dst* index was initiated by Bartels who advocated for the development of an index for monitoring the equatorial ring current (Kertz, 1958, 1964; Mayaud, 1980; Sugiura, 1964). The *Dst* index represents the longitudinally averaged part of the external field measured at the geomagnetic dipole equator at the surface of the Earth (Sugiura, 1964) and is provided by the World Data Center for Geomagnetism, Kyoto. The most often used definition of a geomagnetic storm is an event wherein the minimum of the *Dst* index goes below a critical value, for example, -50 nT or -100 nT (Sugiura & Chapman, 1960). A minimum-*Dst* categorization that is often used to categorize storm intensity is (cf. Table 1 of Loewe & Prolss, 1997) a weak storm has -30 nT $> Dst > -50$ nT, a moderate storm has -50 nT $> Dst > -100$ nT, a strong storm has -100 nT $> Dst > -200$ nT, a severe storm has -200 nT $> Dst > -350$ nT, and a great storm has $Dst < -350$ nT.

Dst is primarily a measure of plasma pressure in the dipolar magnetosphere (Dessler & Parker, 1959; Greenspan & Hamilton, 2000; Schopke, 1966); the *Dst* index is taken to be the strength of the Earth's ring current, but it also reacts to the partial ring current (Liemohn et al., 2001), the cross-tail current (Dubragin et al., 2014; Ohtani et al., 2001; Turner et al., 2000), and the Chapman-Ferraro dayside-magnetopause current (Burton, McPherron, & Russell, 1975; Siscoe, McPherron, & Jordanova, 2005). While the *Dst* index estimates the strength of the storm time ring current, it is a poor proxy for other storm time phenomena. Ionospheric Joule heating during storms is not strongly related to the *Dst* index; it is instead parameterized by high-latitude geomagnetic indices (Baumjohann & Kamide, 1984; Chun et al., 2002). Ionospheric ion outflow is typically parameterized by *Kp* (Welling et al., 2015; Yau, Peterson, & Abe, 2011). Ionospheric models also use either solar wind-driven electric field models or the *Kp* index to evaluate the effects of geomagnetic storms (Garcia et al., 2007; Weimer, 1995). Variations in the thermospheric density (important for atmospheric drag and spacecraft reentry) vary with geomagnetic activity as parameterized by *Kp* (Rhoden, Forbes, & Marcos, 2000; Vallardo & Finkleman, 2014). The energy of auroral particle precipitation is parameterized by *Kp* (Emery et al., 2008; Hardy et al., 1987). *Dst* does not describe the dynamics of the particle populations in the radiation belts (Reeves et al., 2003). None of these quantities are well correlated with *Dst*. The evolution of the hot-plasma ions and electrons in the magnetosphere is well correlated with the *Kp* index (Denton et al., 2015; Korth et al., 1999). The storm time evolution of the cold plasma of the plasmasphere and plumes shows little correlation with *Dst* as it is determined by the time history of the convection electric field, which better correlates with the *Kp* or *AE* index (e.g., Goldstein et al., 2014; Maynard & Chen, 1975; Pierrard et al., 2009).

Dst is also often used to determine whether or not a storm occurred, to define the duration of a storm, and to distinguish between quiet and disturbed geomagnetic conditions. However, *Dst* does not identify all storms. Further, as should be expected, *Dst* is incomplete in describing storm evolution and storm phenomena. The direct relevance of *Dst* to modern space-weather phenomena is weak.

Long-duration high-speed-stream-driven (aka CIR-driven) geomagnetic storms, which lead to the strongest electron-radiation-belt radiation hazards, to the strongest plasma sheet spacecraft-charging hazards, and to the related spacecraft anomalies (Wrenn, 2009; Wrenn, Rodgers, & Ryden, 2002), have weak, short-lived *Dst* signatures (Borovsky & Denton, 2006; McPherron & Weygand, 2006) and are typically not captured in $Dst \leq -50$ nT or $Dst \leq -100$ nT lists of storms (Denton et al., 2009). One reason for this misidentification is




Figure 1. (a) Hourly values of $-Dst$ are binned, and (b) hourly values of the radiation-belt index $F_{e1.2}$ are plotted. In both panels the red curves are for times when $Dst \leq -100$ nT; the dark-green, light-green, and blue curves are for the first, second, and third days of high-speed-stream-driven storms; and the black dashed curve is for all times.

the fact that high-speed-stream-driven storms are characterized by a daylong increase in the solar wind ram pressure at the onset of the storm, which causes intensified Chapman-Ferraro currents to produce a positive perturbation to Dst that offsets the negative perturbation produced by storm convection (Borovsky & Denton, 2010). A second reason is that the ion plasma sheet is hot during a high-speed-stream-driven storm and hot ions drift azimuthally and cannot be injected close to the Earth, which makes them poor fuel for the ring current (Ebihara & Ejiri, 2000; Kozyra & Liemohn, 2003; Liemohn et al., 2001), leading to a weaker negative Dst signature (Lavraud et al., 2006).

A glaring example of an important storm time space-weather phenomena that is not well organized by the Dst index is the evolution of the outer electron radiation belt. In Figure 1a the hourly values of the $-Dst$ index are binned: the red curve is the distribution of $-Dst$ values when $Dst \leq -100$ nT (Dst -defined storms), and the dark-green, light-green, and blue curves are the distributions of $-Dst$ values for the first day, the second day, and the third day, respectively, of 70 high-speed-stream-driven storms from the collection of Borovsky and Denton (2016). The black dashed curve is the distribution of $-Dst$ for all times. Figure 1a indicates that very few of the hours of high-speed-stream-driven storms would be categorized as a storm by the Dst index (less than 1% in Figure 1a). In Figure 1b the multispacecraft 1.2 MeV radiation-belt-intensity index $F_{e1.2}$ (cf. Borovsky & Yakymenko, 2017) is binned for the same events as in Figure 1a: as can be seen, the outer electron radiation belt is weak during the $Dst \leq -100$ nT storms (red) and the radiation belt is strong during high-speed-stream-driven storms, getting stronger every day while Dst remains weak. Statistical parameterizations of the ULF waves responsible for the inward and outward radial transport of radiation-belt particles are usually given in terms of the Kp index (e.g., Brautigam & Albert, 2000; Ozeke et al., 2014). VLF waves that are responsible for acceleration and loss of radiation-belt particles are also usually parameterized by Kp (e.g., Agapitov et al., 2015; Orlova, Shprits, & Spasojevic, 2016; Orlova, Spasojevic, & Shprits, 2014; Shprits, Meredith, & Thorne, 2007; Spasojevic, Shprits, & Orlova, 2015) or by AE/AL (Li et al., 2015; Shprits et al., 2007). The Dst index, however, can serve as a proxy for the adiabatic changes (Kim & Chan, 1997) of electron fluxes driven by the change in the global magnetic field; these changes result in reversible changes in the radiation environment. The behavior of the electron radiation belt is much better organized by an index such as Kp , which measures the strength of magnetospheric convection (Thomsen, 2004). At times of increasing electron-radiation-belt intensities the correlations of the relativistic-electron fluxes with the Kp index are nearly two times stronger than the correlations with

the Dst index (Borovsky, 2017). Similarly for times when the radiation-belt flux is decaying, flux correlations with Kp are much stronger than flux correlations with Dst .

The Dst index is also of limited relevance to other space-weather phenomena. (1) Spacecraft charging hazards are caused by anomalously hot plasma in the magnetosphere, which is more related to Kp and to substorm activity. (2) Geomagnetically induced currents are caused by high-latitude currents, and this hazard is more related to the behavior of high-latitude magnetometer stations. (3) Ionospheric disturbances associated with storm time convection are more related to Kp . And of course the Dst index is irrelevant to the intensity of solar energetic particles at Earth and to solar-flare ionospheric disturbances.

There is a need to develop a methodology (A) to identify all storms or (B) to define and identify the different types of storms. Dst alone is not sufficient for either need.

From a solar wind-driver point of view there are two types of storms: CME-driven (shock, sheath, and ejecta) storms and high-speed-stream-driven (CIR then stream) storms. The widely available K_p index could be used to identify both the CME-driven and the high-speed-stream-driven storms, although a K_p index with a time resolution much faster than the currently available 3 h would be needed to determine storm onset times. The NOAA geomagnetic-storm scale (http://www.swpc.noaa.gov/sites/default/files/images/NOAA_scales.pdf) is based on the instantaneous value of K_p . Focusing specifically on high-speed-stream-driven storms where storm duration impacts the radiation-belt intensity, a running average or time integral of K_p would better serve to identify the occurrence of those storms and a time integral of K_p could describe the evolution of the electron radiation belt. Another way to specifically identify high-speed-stream-driven storms could be via temporal increases in the intensity of the outer electron radiation belt (cf. Figure 1b) that bring the intensity levels to above-average values. A current limitation of the K_p index is that it saturates at $K_p = 9$ and cannot be used to compare the strengths of superstorms for which K_p reaches the maximum allowed value of 9. The discrete values at which K_p is given (e.g., 0, 0+, 1, 1–, and 1+) complicate the systematic analysis of this index. The A_p or the am indices (Mayaud, 1980) may be much more appropriate for gauging storm intensity and for statistical studies.

From a geospace-response point of view there are several types of geospace storms. Focusing on the ring current and the associated low-latitude magnetic disturbances, “geomagnetic storms” as characterized by Dst are important. Focusing on intensifications of relativistic electrons, what might be called “trapped radiation storms” could be characterized by radiation-belt fluxes or by the time integral of a K_p -like index (e.g., Baker, Kanekal, & Blake, 2004; Borovsky & Yakymenko, 2017; Rochel et al., 2016). Focusing on high-latitude currents, “high-latitude electrical storms” could be characterized by dB/dt or the temporal Fourier transforms from ground-based magnetometers (e.g., Marshall et al., 2011; Wintoft, Wik, & Viljanen, 2015). Focusing on ionospheric disturbances associated with convection, “ionospheric storms” could be characterized by f_oF_2 or total electron content (TEC) (e.g., Blagoveshchenskii, 2013; Nishioka et al., 2017). Focusing on spacecraft-charging hazards, “hot-plasma storms” could be characterized by K_p (e.g., Denton et al., 2016; Thomsen, Henderson, & Jordanova, 2013). There are of course overlaps of the various types of geospace storms, just as a thunderstorm can be a rainstorm, a lightning storm, a hailstorm, and/or a windstorm.

From a system-science point of view, a “state vector” composed of several measures would provide a superior indication of the intensity of global geospace activity (Luo, 2010; Vassiliadis, 2006). Storm occurrence and storm severity could be indicated (a) by the nature of the vector, (b) by the magnitude of the vector, or (c) by the dot product of the vector with some indicator of storm properties. Components of such a state vector might be measurements of critical magnetospheric properties such as the amount of open flux, the strength of convection, the intensity of the ring current ions, the density of the cold plasma population, the intensity of the electron radiation belt, the substorm-occurrence rate, the intensity of the substorm-injected energetic-particle population, the plasma sheet density and temperature, indices of ULF wave activity, ionospheric global TEC, and f_oF_2 indices. Geomagnetic indices such as Dst , K_p , AL , and PCI should also be components of the state vector.

An approach could be to specifically catalog different types of storms separately using different definitions and different criteria. With the state-vector approach it might involve taking the dot product of the magnetospheric state vector with different reference vectors to highlight properties that are characteristic of the different types of storms.

The Dst index alone is not an appropriate indicator for identifying all geospace storms, for the comparison of the evolution during different storms, for performing superposed epoch analysis, or for determining the onset and end of a storm.

The physical processes operating during storms are much better understood now than they were when von Humbolt originally defined a magnetic storm and when the indices were first created by Bartels, Kertz, and Sugiura. Many studies choose Dst as a proxy for storms simply because of its continuous scale, its high cadence, and its simple interpretation; however, as argued in this commentary such use of this index may often be inappropriate or not optimal for identification of geomagnetic storms and representation of storms and their effects. We as a community should consider a better set of criteria.

Acknowledgments

The authors thank Mick Denton, Mike Henderson, Herman Luhr, Juergen Matzka, Claudia Stolle, and Michelle Thomsen for helpful conversations. Work at the Space Science Institute was supported by the NASA Heliophysics LWS TRT program via grants NNX14AN90G and NNX16AB75G, by the NSF GEM Program via award AGS-1502947, by the NSF Solar-Terrestrial Program via grant AGS-12GG13659, and by the NASA Heliophysics Guest Investigator Program via grant NNX14AC15G. Work at UCLA was supported by NASA HSR NNX15AI94G and HSR NNX16AG78G and by subcontract 1554053 from the University of Colorado under NASA grant NNX16AF91G. Work at GDZ was supported by the European Union's Horizon 2020 research and innovation program under grant agreement 637302 and also by SFB 1294. Data from the plots can be obtained from J.E.B.

References

- Agapitov, O. V., Artemyev, A. V., Mourenas, D., Mozer, F. S., & Krasnoselskikh, V. (2015). Empirical model of lower band chorus wave distribution in the outer radiation belt. *Journal of Geophysical Research: Space Physics*, 120, 10,425–10,442. <https://doi.org/10.1002/2015JA021829>
- Baker, D. N., Kanekal, S. G., & Blake, J. B. (2004). Characterizing the Earth's outer Van Allen zone using a radiation belt content index. *Space Weather*, 2, S02003. <https://doi.org/10.1029/2003SW000026>
- Baumjohann, W., & Kamide, Y. (1984). Hemispherical Joule heating and the AE indices. *Journal of Geophysical Research*, 89, 383–388. <https://doi.org/10.1029/JA089iA01p00383>
- Blagoveshchenskii, D. V. (2013). Effect of geomagnetic storms (substorms) on the ionosphere: 1. A review. *Geomagnetism and Aeronomy*, 53(3), 275–290. <https://doi.org/10.1134/S0016793213030031>
- Borovsky, J. E. (2017). Time-integral correlations of multiple variables with the relativistic-electron flux at geosynchronous orbit: The strong roles of the substorm-injected electrons and the ion plasma sheet. *Journal of Geophysical Research: Space Physics*, 122. <https://doi.org/10.1002/2017JA024476>
- Borovsky, J. E., & Denton, M. H. (2006). The differences between CME-driven storms and CIR-driven storms. *Journal of Geophysical Research*, 111, A07S08. <https://doi.org/10.1029/2005JA011447>
- Borovsky, J. E., & Denton, M. H. (2010). The magnetic field at geosynchronous orbit during high-speed-stream-driven storms: Connections to the solar wind, the plasma sheet, and the outer electron radiation belt. *Journal of Geophysical Research*, 115, A08217. <https://doi.org/10.1029/2009JA015116>
- Borovsky, J. E., & Denton, M. H. (2016). Compressional perturbations of the dayside magnetosphere during high-speed-stream-driven geomagnetic storms. *Journal of Geophysical Research*, 121, 4569–4589. <https://doi.org/10.1002/2015JA022136>
- Borovsky, J. E., & Yakymenko, K. (2017). Systems science of the magnetosphere: Creating indices of substorm activity, of the substorm-injected electron population, and of the electron radiation belt. *Journal of Geophysical Research: Space Physics*, 122. <https://doi.org/10.1002/2017JA024250>
- Brautigam, D. H., & Albert, J. M. (2000). Radial diffusion analysis of outer radiation belt electrons during the October 9, 1990, magnetic storm. *Journal of Geophysical Research*, 105(A1), 291–309. <https://doi.org/10.1029/1999JA900344>
- Burton, R. K., McPherron, R. L., & Russell, C. T. (1975). An empirical relationship between interplanetary conditions and Dst. *Journal of Geophysical Research*, 80(31), 4204–4214. <https://doi.org/10.1029/JA080i031p04204>
- Chun, F. K., Knipp, D. J., McHarg, M. G., Lacey, J. R., Lu, G., & Emery, B. A. (2002). Joule heating patterns as a function of polar cap index. *Journal of Geophysical Research*, 107(A7), 1119. <https://doi.org/10.1029/2001JA000246>
- Denton, M. H., Borovsky, J. E., Horne, R. B., McPherron, R. L., Morley, S. K., & Tsurutani, B. T. (2009). Introduction to Special Issue on high speed solar wind streams and geospace interactions (HSS-GI). *Journal of Atmospheric and Solar - Terrestrial Physics*, 71(10–11), 1011–1013. <https://doi.org/10.1016/j.jastp.2008.09.019>
- Denton, M. H., Henderson, M. G., Jordanova, V. K., Thomsen, M. F., Borovsky, J. E., Woodroffe, J., ... Pitchford, D. (2016). An improved model of electron and ion fluxes at geosynchronous orbit based on upstream solar wind conditions. *Space Weather*, 14, 511–523. <https://doi.org/10.1002/2016SW001409>
- Denton, M. H., Thomsen, M. F., Jordanova, V. K., Henderson, M. G., Borovsky, J. E., Denton, J. S., ... Hartley, D. P. (2015). An empirical model of electron and ion fluxes derived from observations at geosynchronous orbit. *Space Weather*, 13, 233–249. <https://doi.org/10.1002/2015SW001168>
- Dessler, A. J., & Parker, E. N. (1959). Hydromagnetic theory of geomagnetic storms. *Journal of Geophysical Research*, 64(12), 2239–2252. <https://doi.org/10.1029/JZ064i012p02239>
- Dubyagin, S., Ganushkina, N., Kubyshkina, M., & Liemohn, M. (2014). Contribution from different current systems to SYM and ASY midlatitude indices. *Journal of Geophysical Research*, 119, 7243.
- Ebihara, Y., & Ejiri, M. (2000). Simulation study on fundamental properties of the storm-time ring current. *Journal of Geophysical Research*, 105(A7), 15,843–15,859. <https://doi.org/10.1029/1999JA900493>
- Emery, B. A., Coumans, V., Evans, D. S., Germany, G. A., Greer, M. S., Holeman, E., ... Xu, W. (2008). Seasonal Kp, solar wind, and solar flux variations in long-term single-pass satellite estimates of electron and ion auroral hemispheric power. *Journal of Geophysical Research*, 113, A06311. <https://doi.org/10.1029/2007JA012866>
- Garcia, R. R., Marsh, D. R., Kinnison, D. E., Boville, B. A., & Sassi, F. (2007). Simulation of secular trends in the middle atmosphere, 1950–2003. *Journal of Geophysical Research*, 112, D09301. <https://doi.org/10.1029/2006JD007485>
- Goldstein, J., De Pascuale, S., Kletzing, C., Kurth, W., Genestreti, K. J., Skoug, R. M., ... Spence, H. (2014). Simulation of Van Allen Probes plasmopause encounters. *Journal of Geophysical Research: Space Physics*, 119, 7464–7484. <https://doi.org/10.1002/2014JA020252>
- Greenspan, M. E., & Hamilton, D. C. (2000). A test of the Dessler-Parker-Scokpe relation during magnetic storms. *Journal of Geophysical Research*, 105(A3), 5419–5430. <https://doi.org/10.1029/1999JA000284>
- Hardy, D. A., Gussenhoven, M. S., Raistrick, R., & McNeil, W. J. (1987). Statistical and functional representations of the pattern of auroral energy flux, number flux, and conductivity. *Journal of Geophysical Research*, 92(A11), 12,275–12,294. <https://doi.org/10.1029/JA092iA11p12275>
- Kertz, W. (1958). Ein neues Mass für die Feldstärke des erdmagnetischen aquatorialen Ringstroms. *Abhandlungen der Akademie der Wissenschaften Göttingen der mathemat.-physikalischen*, 2, 83.
- Kertz, W. (1964). Ring current variations during the IGY. In *Annals of the International Geophysical Year* (Vol. 35, pp. 49–61). London: Pergamon Press.
- Kim, H.-J., & Chan, A. A. (1997). Fully adiabatic changes in storm time relativistic electron fluxes. *Journal of Geophysical Research*, 102(A10), 22,107–22,116. <https://doi.org/10.1029/97JA01814>
- Korth, H., Thomsen, M. F., Borovsky, J. E., & McComas, D. J. (1999). Plasma sheet access to geosynchronous orbit. *Journal of Geophysical Research*, 104(A11), 25,047–25,061. <https://doi.org/10.1029/1999JA900292>
- Kozyra, J. U., & Liemohn, M. W. (2003). Ring current energy input and decay. *Space Science Reviews*, 109(1–4), 105–131. <https://doi.org/10.1023/B:SPAC.0000007516.10433.ad>
- Lavraud, B., Thomsen, M. F., Borovsky, J. E., Denton, M. H., & Pulkkinen, T. I. (2006). Magnetosphere preconditioning under northward IMF: Evidence from the study of coronal mass ejection and corotating interaction region geoeffectiveness. *Journal of Geophysical Research*, 111, A09208. <https://doi.org/10.1029/2005JA011566>
- Li, W., Ma, Q., Thorne, R. M., Bortnik, J., Kletzing, C. A., Kurth, W. S., ... Nishimura, Y. (2015). Statistical properties of plasmaspheric hiss derived from Van Allen Probes data and their effects on radiation belt electron dynamics. *Journal of Geophysical Research: Space Physics*, 120, 3393–3405. <https://doi.org/10.1002/2015JA021048>

- Liemohn, M. W., Kozyra, J. U., Thomsen, M. F., Roeder, J. L., Lu, G., Borovsky, J. E., & Cayton, T. E. (2001). Dominant role of the asymmetric ring current in producing stormtime *Dst**. *Journal of Geophysical Research*, 106(A6), 10,883–10,904. <https://doi.org/10.1029/2000JA000326>
- Loewe, C. A., & Prolss, G. W. (1997). Classification of mean behavior of magnetic storms. *Journal of Geophysical Research*, 102(A7), 14,209–14,213. <https://doi.org/10.1029/96JA04020>
- Luo, A. C. J. (Ed.) (2010). *Dynamical systems*. New York: Springer. <https://doi.org/10.1007/978-1-4419-5754-2>
- Malin, S. R. C., & Barraclough, D. R. (1991). *Quarterly Journal of the Royal Astronomical Society*, 32, 279.
- Marshall, R. A., Smith, E. A., Francis, M. J., Waters, C. L., & Sciffer, M. D. (2011). A preliminary risk assessment of the Australian region power network to space weather. *Space Weather*, 9, S10004. <https://doi.org/10.1029/2011SW000685>
- Mayaud, P. N. (1980). *Derivation, meaning, and use of geomagnetic indices* (Sect. 5.1.3, Chap. 8). Washington, DC: American Geophysical Union. <https://doi.org/10.1029/GM022>
- Maynard, N., & Chen, A. (1975). Isolated cold plasma regions: Observations and their relation to possible production mechanisms. *Journal of Geophysical Research*, 80(7), 1009–1013. <https://doi.org/10.1029/JA080i007p01009>
- McPherron, R. L., & Weygand, J. (2006). The solar wind and geomagnetic activity as a function of time relative to corotating interaction regions. In B. Tsurutani, et al. (Eds.), *Recurrent magnetic storms* (pp. 125–137). Washington, DC: American Geophysical Union. <https://doi.org/10.1029/167GM12>
- Nishioka, M., Tsugawa, T., Jin, H., & Ishii, M. (2017). A new ionospheric storm scale based on TEC and F_oF_2 statistics. *Space Weather*, 15, 228–239. <https://doi.org/10.1002/2016SW001536>
- Ohtani, S., Nose, M., Rostoker, G., Singer, H., Lui, A. T. Y., & Nakamura, M. (2001). Storm-substorm relationship: Contribution of the tail current to *Dst*. *Journal of Geophysical Research*, 106(A10), 21,199–21,209. <https://doi.org/10.1029/2000JA000400>
- Orlova, K., Shprits, Y., & Spasojevic, M. (2016). New global loss model of energetic and relativistic electrons based on Van Allen Probes measurements. *Journal of Geophysical Research: Space Physics*, 121, 1308–1314. <https://doi.org/10.1002/2015JA021878>
- Orlova, K., Spasojevic, M., & Shprits, Y. (2014). Activity-dependent global model of electron loss inside the plasmasphere. *Geophysical Research Letters*, 41, 3744–3751. <https://doi.org/10.1002/2014GL060100>
- Ozeke, L. G., Mann, I. R., Murphy, K. R., Rae, I. J., & Milling, D. K. (2014). Analytic expressions for ULF wave radiation belt radial diffusion coefficients. *Journal of Geophysical Research: Space Physics*, 119, 1587–1605. <https://doi.org/10.1002/2013JA019204>
- Pierrard, V., Goldstein, J., André, N., Jordanova, V. K., Kotova, G. A., Lemaire, J. F., ... Matsui, H. (2009). Recent progress in physics-based models of the plasmasphere. *Space Science Reviews*, 145(1–2), 193–229. <https://doi.org/10.1007/s11214-008-9480-7>
- Reeves, G. D., McAdams, K. L., Friedel, R. H. W., & O'Brien, T. P. (2003). Acceleration and loss of relativistic electrons during geomagnetic storms. *Geophysical Research Letters*, 30(10), 1529. <https://doi.org/10.1029/2002GL016513>
- Rhoden, E. A., Forbes, J. M., & Marcos, F. A. (2000). The influence of geomagnetic and solar variabilities on lower thermospheric densities. *Journal of Atmospheric and Solar - Terrestrial Physics*, 62(11), 999–1013. [https://doi.org/10.1016/S1364-6826\(00\)00066-3](https://doi.org/10.1016/S1364-6826(00)00066-3)
- Rochel, S., Boscher, D., Benacquista, R., & Roussel, J. F. (2016). A radiation belt disturbance study from the space weather point of view. *Acta Astronautica*, 128, 650–656. <https://doi.org/10.1016/j.actaastro.2016.07.012>
- Scopke, N. (1966). A general relation between the energy of trapped particles and the disturbance field near the Earth. *Journal of Geophysical Research*, 71(13), 3125–3130. <https://doi.org/10.1029/JZ071i013p03125>
- Shprits, Y. Y., Meredith, N. P., & Thorne, R. M. (2007). Parameterization of radiation belt electron loss timescales due to interactions with chorus waves. *Geophysical Research Letters*, 34, L11110. <https://doi.org/10.1029/2006GL029050>
- Siscoe, G. L., McPherron, R. L., & Jordanova, V. K. (2005). Diminished contribution of ram pressure to *Dst* during magnetic storms. *Journal of Geophysical Research*, 110, A12227. <https://doi.org/10.1029/2005JA011120>
- Spasojevic, M., Shprits, Y. Y., & Orlova, K. (2015). Global empirical models of plasmaspheric hiss using Van Allen Probes. *Journal of Geophysical Research: Space Physics*, 120, 10,370–10,383. <https://doi.org/10.1002/2015JA021803>
- Sugiura, M. (1964). Hourly values of equatorial *Dst* for the IGY. In *Annals of the International Geophysical Year* (Vol. 35, pp. 7–45). Oxford: Pergamon Press.
- Sugiura, M., & Chapman, S. (1960). The average morphology of geomagnetic storms with sudden commencement. In *Abhandlungen der Akademie der Wissenschaften zu Göttingen* (pp. 1–53). Göttingen: Göttingen Math. Phys. Kl., Sonderheft Nr.4.
- Thomsen, M. F. (2004). Why *Kp* is such a good measure of magnetospheric convection. *Space Weather*, 2, S11044. <https://doi.org/10.1029/2004SW000089>
- Thomsen, M. F., Henderson, M. G., & Jordanova, V. (2013). Statistical properties of the surface-charging environment at geosynchronous orbit. *Space Weather*, 11, 237–244. <https://doi.org/10.1002/swe.20049>
- Turner, N. E., Baker, D. N., Pulkkinen, T. I., & McPherron, R. L. (2000). Evaluation of the tail current contribution to *Dst*. *Journal of Geophysical Research*, 105(A3), 5431–5439. <https://doi.org/10.1029/1999JA000248>
- Vallardo, D. A., & Finkleman, D. (2014). A critical assessment of satellite drag and atmospheric density modeling. *Acta Astronautica*, 95, 141–165. <https://doi.org/10.1016/j.actaastro.2013.10.005>
- Vassiliadis, D. (2006). Systems theory for geospace plasma dynamics. *Reviews of Geophysics*, 44, RG2002. <https://doi.org/10.1029/2004RG000161>
- Weimer, D. R. (1995). Models of high-latitude electric potentials derived with a least error fit of spherical harmonic coefficients. *Journal of Geophysical Research*, 100(A10), 19,595–19,607. <https://doi.org/10.1029/95JA01755>
- Welling, D. T., Andre, M., Dandouras, I., Delcourt, D., Fazakerley, A., Fontaine, D., ... Yau, A. (2015). The Earth: Plasma sources, losses, and transport processes. *Space Science Reviews*, 192(1–4), 145–208. <https://doi.org/10.1007/s11214-015-0187-2>
- Wintoft, P., Wik, M., & Viljanen, A. (2015). Solar wind driven empirical forecast models of the time derivative of the ground magnetic field. *Journal of Space Weather and Space Climate*, 5, A7. <https://doi.org/10.1051/swsc/2015008>
- Wrenn, G. L. (2009). Chronology of “killer” electron: Solar cycles 22 and 23. *Journal of Atmospheric and Solar - Terrestrial Physics*, 71(10–11), 1210–1218. <https://doi.org/10.1016/j.jastp.2008.08.002>
- Wrenn, G. L., Rodgers, D. J., & Ryden, K. A. (2002). A solar cycle of spacecraft anomalies due to internal charging. *Annales de Geophysique*, 20(7), 953–956. <https://doi.org/10.5194/angeo-20-953-2002>
- Yau, A. W., Peterson, W. K., & Abe, T. (2011). Influences of the ionosphere, thermosphere and magnetosphere on ion outflows. In W. Liu & M. Fujimoto (Eds.), *The dynamic magnetosphere* (pp. 283–314). Dordrecht, Netherlands: Springer Science+Business Media. https://doi.org/10.1007/978-94-007-0501-2_16